

Asymptotic expansions for ratios of products of gamma functions

Wolfgang Bühring

Physikalisches Institut, Universität Heidelberg, Philosophenweg 12
69120 Heidelberg, Germany

buehring@physi.uni-heidelberg.de

Abstract

An asymptotic expansion for a ratio of products of gamma functions is derived.

2000 Mathematics Subject Classification: Primary 33B15; Secondary 33C20

Keywords and phrases: Gamma function, generalized hypergeometric functions

1 Introduction

An asymptotic expansion for a ratio of products of gamma functions has recently been found [2], which, with

$$s_1 = b_1 - a_1 - a_2, \quad (1)$$

may be written

$$\frac{\Gamma(a_1 + n)\Gamma(a_2 + n)}{\Gamma(b_1 + n)\Gamma(-s_1 + n)} = 1 + \sum_{m=1}^M \frac{(s_1 + a_1)_m (s_1 + a_2)_m}{(1)_m (1 + s_1 - n)_m} + O(n^{-M-1}) \quad (2)$$

as $n \rightarrow \infty$. Here use is made of the Pochhammer symbol

$$(x)_n = x(x+1) \cdots (x+n-1) = \Gamma(x+n)/\Gamma(x).$$

The special case when $b_1 = 1$ of this formula (2) had been stated earlier by Dingle [3], and there were proofs by Paris [8] and Olver [6][7].

The proof of (2) is based on the formula for the analytic continuation near unit argument of the Gaussian hypergeometric function. For the more general hypergeometric functions

$${}_{p+1}F_p \left(\begin{matrix} a_1, a_2, \dots, a_{p+1} \\ b_1, \dots, b_p \end{matrix} \middle| z \right) = \sum_{n=0}^{\infty} \frac{(a_1)_n (a_2)_n \cdots (a_{p+1})_n}{(b_1)_n \cdots (b_p)_n (1)_n} z^n, \quad (|z| < 1), \quad (3)$$

the analytic continuation near $z = 1$ is known too, and this raises the question as to whether a sufficiently simple asymptotic expansion can be derived in a similar way for a ratio of products of more gamma function factors. This is indeed the case, and it is the purpose of this work to present such an expansion.

2 Derivation of the asymptotic expansion

The analytic continuation of the hypergeometric function near unit argument may be written [1]

$$\begin{aligned} & \frac{\Gamma(a_1)\Gamma(a_2) \cdots \Gamma(a_{p+1})}{\Gamma(b_1) \cdots \Gamma(b_p)} {}_{p+1}F_p \left(\begin{matrix} a_1, a_2, \dots, a_{p+1} \\ b_1, \dots, b_p \end{matrix} \middle| z \right) \\ &= \sum_{m=0}^{\infty} g_m(0) (1-z)^m + (1-z)^{s_p} \sum_{m=0}^{\infty} g_m(s_p) (1-z)^m, \end{aligned} \quad (4)$$

where

$$s_p = b_1 + \cdots + b_p - a_1 - a_2 - \cdots - a_{p+1} \quad (5)$$

and the coefficients g_m are known. While the $g_m(0)$ are not needed for the present purpose, the $g_m(s_p)$ are [1]

$$\begin{aligned} g_m(s_p) &= (-1)^m \frac{(a_1 + s_p)_m (a_2 + s_p)_m \cdots \Gamma(-s_p - m)}{(1)_m} \\ &\times \sum_{k=0}^m \frac{(-m)_k}{(a_1 + s_p)_k (a_2 + s_p)_k} A_k^{(p)}, \end{aligned} \quad (6)$$

where the coefficients $A_k^{(p)}$ will be shown below.

The left-hand side L of (4) is

$$L = \sum_{n=0}^{\infty} \frac{\Gamma(a_1 + n)\Gamma(a_2 + n) \cdots \Gamma(a_{p+1} + n)}{\Gamma(b_1 + n) \cdots \Gamma(b_p + n)\Gamma(1 + n)} z^n. \quad (7)$$

The asymptotic behaviour, as $n \rightarrow \infty$, of the coefficients of this power series is governed [4][5][10] by the terms R on the right-hand side which, at $z = 1$, are singular,

$$R = \sum_{m=0}^{\infty} g_m(s_p)(1 - z)^{s_p+m}. \quad (8)$$

Expanded by means of the binomial theorem in its hypergeometric-series-form, this is

$$R = \sum_{m=0}^{\infty} g_m(s_p) \sum_{n=0}^{\infty} \frac{(-s_p - m)_n}{\Gamma(1 + n)} z^n. \quad (9)$$

Interchanging the order of summation (and making use of the reflection formula of the gamma function) we may get

$$R = \sum_{n=0}^{\infty} \frac{\Gamma(-s_p + n)}{\Gamma(1 + n)} \sum_{m=0}^{\infty} (-1)^m g_m(s_p) \frac{1}{\Gamma(-s_p - m)(1 + s_p - n)_m} z^n. \quad (10)$$

Comparison of the coefficients of the two power series for R and L , which asymptotically, as $n \rightarrow \infty$, should agree, then leads to

$$\begin{aligned} & \frac{\Gamma(a_1 + n)\Gamma(a_2 + n) \cdots \Gamma(a_{p+1} + n)}{\Gamma(b_1 + n) \cdots \Gamma(b_p + n)\Gamma(-s_p + n)} \\ & \sim \sum_{m=0}^{\infty} (-1)^m g_m(s_p) \frac{1}{\Gamma(-s_p - m)(1 + s_p - n)_m}. \end{aligned} \quad (11)$$

Inserting g_m from (6) and keeping the first $M + 1$ terms of the asymptotic series, we get

Theorem 1

$$\begin{aligned} & \frac{\Gamma(a_1 + n)\Gamma(a_2 + n) \cdots \Gamma(a_{p+1} + n)}{\Gamma(b_1 + n) \cdots \Gamma(b_p + n)\Gamma(-s_p + n)} = 1 \\ & + \sum_{m=1}^M \frac{(a_1 + s_p)_m (a_2 + s_p)_m}{(1)_m (1 + s_p - n)_m} \sum_{k=0}^m \frac{(-m)_k}{(a_1 + s_p)_k (a_2 + s_p)_k} A_k^{(p)} + O(n^{-M-1}) \end{aligned} \quad (12)$$

as $n \rightarrow \infty$, where $s_p = b_1 + \cdots + b_p - a_1 - a_2 - \cdots - a_{p+1}$.

The simple formula (2) above corresponding to $p = 1$ can be recovered from this theorem if we define $A_0^{(1)} = 1$, $A_k^{(1)} = 0$ for $k > 0$, so that then the sum over k is equal to 1 and disappears. The coefficients for larger p can be found in [1], but a few of them are here displayed again for convenience:

$$A_k^{(2)} = \frac{(b_2 - a_3)_k (b_1 - a_3)_k}{k!}, \quad (13)$$

$$A_k^{(3)} = \sum_{k_2=0}^k \frac{(b_3 + b_2 - a_4 - a_3 + k_2)_{k-k_2} (b_1 - a_3)_{k-k_2} (b_3 - a_4)_{k_2} (b_2 - a_4)_{k_2}}{(k - k_2)! k_2!}, \quad (14)$$

$$\begin{aligned} A_k^{(4)} = & \sum_{k_2=0}^k \frac{(b_4 + b_3 + b_2 - a_5 - a_4 - a_3 + k_2)_{k-k_2} (b_1 - a_3)_{k-k_2}}{(k - k_2)!} \\ & \times \sum_{k_3=0}^{k_2} \frac{(b_4 + b_3 - a_5 - a_4 + k_3)_{k_2-k_3} (b_2 - a_4)_{k_2-k_3}}{(k_2 - k_3)!} \\ & \times \frac{(b_4 - a_5)_{k_3} (b_3 - a_5)_{k_3}}{k_3!}. \end{aligned} \quad (15)$$

For $p = 3, 4, \dots$, several other representations are possible [1], such like

$$A_k^{(3)} = \frac{(b_3 + b_2 - a_4 - a_3)_k (b_1 - a_3)_k}{k!} \quad (16)$$

$$\times {}_3F_2 \left(\begin{matrix} b_3 - a_4, b_2 - a_4, -k \\ b_3 + b_2 - a_4 - a_3, 1 + a_3 - b_1 - k \end{matrix} \middle| 1 \right)$$

or

$$A_k^{(3)} = \frac{(b_1 + b_3 - a_3 - a_4)_k (b_2 + b_3 - a_3 - a_4)_k}{k!} \quad (17)$$

$$\times {}_3F_2 \left(\begin{matrix} b_3 - a_3, b_3 - a_4, -k \\ b_1 + b_3 - a_3 - a_4, b_2 + b_3 - a_3 - a_4 \end{matrix} \middle| 1 \right).$$

For $p = 2$, the formula (12) may be written simply as

$$\frac{\Gamma(a_1 + n) \Gamma(a_2 + n) \Gamma(a_3 + n)}{\Gamma(b_1 + n) \Gamma(b_2 + n) \Gamma(-s_2 + n)} = 1 \quad (18)$$

$$+ \sum_{m=1}^M \frac{(a_1 + s_2)_m (a_2 + s_2)_m}{(1)_m (1 + s_2 - n)_m} {}_3F_2 \left(\begin{matrix} b_2 - a_3, b_1 - a_3, -m \\ a_1 + s_2, a_2 + s_2 \end{matrix} \middle| 1 \right) + O(n^{-M-1}),$$

where $s_2 = b_1 + b_2 - a_1 - a_2 - a_3$.

3 Additional comments

The derivation of the theorem is based on the continuation formula (4) which holds, as it stands, only if s_p is not equal to an integer. Nevertheless, the theorem is valid without such a restriction. This can be verified if the derivation is repeated starting from any of the continuation formulas for the exceptional cases [1]. Instead of or in addition to the binomial theorem, the expansion

$$(1 - z)^m \ln(1 - z) = \sum_{n=1}^{\infty} c_n z^n, \quad (19)$$

is then needed for integer $m \geq 0$, where

$$c_n = -\frac{1}{n}(-1)^m \frac{\Gamma(1+m)\Gamma(n-m)}{\Gamma(n)} \quad (20)$$

for $n > m$, while the coefficients are not needed here for $n \leq m$.

The theorem has been proved here for any sufficiently large positive integer n only. On the basis of the discussion in [2], it can be suspected that the theorem may be theoretically valid (although less useful) in the larger domain of the complex n -half-plane $\operatorname{Re}(s_p + a_1 + a_2 - 1 + n) \geq 0$.

Expansions for ratios of even more general products of gamma functions are treated in a recent monograph by Paris and Kaminski [9].

References

- [1] W. Bühring, *Generalized hypergeometric functions at unit argument*, Proc. Amer Math. Soc. **114** (1992), 145–153.
- [2] W. Bühring, *An asymptotic expansion for a ratio of products of gamma functions*, Internat. J. Math. & Math. Sci. **24** (2000), 504–510.
- [3] R. B. Dingle, *Asymptotic Expansions: Their Derivation and Interpretation* (Academic Press, London, 1973).
- [4] P. Flajolet and A. Odlyzko, *Singularity analysis of generating functions*, SIAM J. Discrete Math. **3** (1990), 216–240.
- [5] F. W. J. Olver, *Asymptotics and Special Functions* (Academic Press, New York, 1974).

- [6] F. W. J. Olver, *Asymptotic expansions of the coefficients in asymptotic series solutions of linear differential equations*, Methods and Applications of Analysis **1** (1994), 1–13.
- [7] F. W. J. Olver, *On an asymptotic expansion of a ratio of gamma functions*, Proc. Royal Irish Acad. **A95** (1995), 5–9.
- [8] R. B. Paris, *Smoothing of the Stokes phenomenon using Mellin-Barnes integrals*, J. Comput. Appl. Math. **41** (1992), 117–133.
- [9] R. B. Paris and D. Kaminski, *Asymptotics and Mellin-Barnes integrals*, Cambridge University Press, Cambridge, 2001.
- [10] R. Schäfke and D. Schmidt, *The connection problem for general linear ordinary differential equations at two regular singular points with applications in the theory of special functions*, SIAM J. Math. Anal. **11** (1980), 848–862.